



Published in final edited form as:

Anticancer Res. 2010 February ; 30(2): 319–325.

Curcumin Targets FOLFOX-surviving Colon Cancer Cells via Inhibition of EGFRs and IGF-1R

Bhaumik B. Patel^{1,2}, Deepshika Gupta², Althea A. Elliott¹, Vivek Sengupta², Yingjie Yu^{1,3}, and Adhip P. N. Majumdar^{1,2,3}

¹Department of Veterans Affairs Medical Center, Wayne State University, Detroit, MI 48201, U.S.A.

²Karmanos Cancer Center, Wayne State University, Detroit, MI 48201, U.S.A.

³Department of Internal Medicine, Wayne State University, Detroit, MI 48201, U.S.A.

Abstract

Curcumin (diferuloylmethane), which has no discernible toxicity, inhibits initiation, promotion and progression of carcinogenesis. 5-Fluorouracil (5-FU) or 5-FU plus oxaliplatin (FOLFOX) remains the backbone of colorectal cancer chemotherapeutics, but produces an incomplete response resulting in survival of cells (chemo-surviving cells) that may lead to cancer recurrence. The present investigation was, therefore, undertaken to examine whether addition of curcumin to FOLFOX is a superior therapeutic strategy for chemo-surviving cells. Forty-eight-hour treatment of colon cancer HCT-116 and HT-29 cells with FOLFOX resulted in 60–70% survival, accompanied by a marked activation of insulin like growth factor-1 receptor (IGF-1R) and minor to moderate increase in epidermal growth factor receptor (EGFR), v-erb-b2 erythroblastic leukemia viral oncogene homolog 2 (HER-2) as well as v-akt murine thymoma viral oncogene homolog 1 (AKT), cyclooxygenase-2 (COX-2) and cyclin-D1. However, inclusion of curcumin to continued FOLFOX treatment for another 48 h greatly reduced the survival of these cells, accompanied by a concomitant reduction in activation of EGFR, HER-2, IGF-1R and AKT, as well as expression of COX-2 and cyclin-D1. More importantly, EGFR tyrosine kinase inhibitor gefitinib or attenuation of IGF-1R expression by the corresponding si-RNA caused a 30–60% growth inhibition of chemo-surviving HCT-116 cells. However, curcumin alone was found to be more effective than both gefitinib and IGF-1R si-RNA mediated growth inhibition of chemo-surviving HCT-116 cells and addition of FOLFOX to curcumin did not increase the growth inhibitory effect of curcumin. Our data suggest that inclusion of curcumin in conventional chemotherapeutic regimens could be an effective strategy to prevent the emergence of chemoresistant colon cancer cells.

Keywords

Chemo-surviving cells; curcumin; colon cancer; EGFRs; IGF-1R

Despite recent incorporation of novel cytotoxic chemotherapy and biological agents in the treatment of metastatic colorectal cancer, the outcome remains poor (1, 2). Currently, 5-fluorouracil (5-FU) or 5-FU plus oxaliplatin (FOLFOX) remains the backbone of colorectal cancer chemotherapeutics. Although chemotherapeutic regimens containing FOLFOX

produce a response in the majority of the cases, virtually all the responses are incomplete and emergence of resistance, with subsequent recurrence of the cancer, is universal.

Accumulating evidence suggests that the development and progression of many malignancies, including colorectal cancer, are associated with constitutive activation of multiple signaling pathways that promote proliferation, inhibit apoptosis and induce metastasis (3). A large body of evidence suggests that EGFR and/or its family members, specifically ErbB-2/HER-2 and ErbB-3/HER-3 (collectively referred to as EGFRs) play a crucial role in regulating several pathways that affect tumor cell survival, angiogenesis, motility and invasiveness (4–6). Insulin-like growth factor (IGF)/IGF-1R system has also been implicated in the development and progression of colorectal cancer (7, 8). In addition, EGFRs have been shown to mediate resistance to chemotherapeutic agents (9–11). Therefore, agent(s) that would target EGFRs and IGF-1R are likely to affect multiple aspects of tumor survival and chemoresistance.

Curcumin (diferuloylmethane), the major active ingredient of turmeric (*Curcuma longa*) with no discernable toxicity, has been shown to inhibit the growth of transformed cells and colon carcinogenesis at the initiation, promotion and progression stages in carcinogen-induced rodent models (12–16). Curcumin has also been shown to prevent the development of adenomas in the intestinal tract of Min^{+/-} mice, a model of human familial adenomatous polyposis (17). In a phase I clinical trial, curcumin has been found to be effective in inhibiting the growth of a variety of tumors (18). Recently, we reported that curcumin synergizes with FOLFOX by inhibiting all EGFRs as well as IGF-1R in colon cancer cells (19).

However, it remains to be determined whether curcumin would be effective against chemo-surviving cells, the cells that survive acute chemotherapy insult. The current *in vitro* study was undertaken to examine the effect of curcumin on the growth of chemo-surviving colon cancer cells (cells that survived 48-h treatment with FOLFOX). We have also examined the possible mechanism of growth inhibition focusing on the regulation of growth factor receptors pathways, specifically EGFRs and IGF-1R which are considered to play an important role in the progression of colorectal cancer.

Materials and Methods

Cell lines and cell cultures

Human colon cancer HCT-116 and HT-29 cells were obtained from the American Type Culture Collection (ATCC, Rockville, MD, USA). The cells were maintained in Dulbecco's modified Eagle's medium (DMEM) in tissue culture flasks in a humidified incubator at 37°C in an atmosphere of 95% air and 5% CO₂. The medium was supplemented with 10% fetal bovine serum (FBS) and 1% antibiotic/anti-mycotic and was changed three times a week. Cells were passaged using trypsin/EDTA. Curcumin was purchased from Sigma Chemical Co. (St. Louis, MO, USA) and dissolved in 100% ethanol and subsequently diluted with the tissue culture medium.

Growth inhibition assay

Inhibition of cell growth in response to curcumin and/or FOLFOX was assessed by 3-(4,5-dimethylthiazol-2yl)-2,5-diphenyltetrazolium bromide (MTT) assay as described previously (15, 24). Briefly, cells were dispersed by trypsin-EDTA treatment and 2.5×10⁴ cells/ml, re-suspended in DMEM containing 10% FBS and seeded into 96-wells culture plates with 6 replicates. After 24 h of plating, incubation was continued for another 48 h in the absence (control) or presence of FOLFOX alone and in combination with 25 μM or 50 μM of curcumin. At the end of the 48 h incubation period, the reaction was terminated by adding

20 μ l of 5 mg/ml stock of MTT to each well. The reaction was allowed to proceed for 3–4 hours at 37°C. The formazan crystals were then dissolved by adding 0.1 ml of dimethyl sulfoxide (DMSO). The intensity of the color developed, which is the reflection of number of live cells, was measured at a wavelength of 570 nm. All values were compared to the corresponding controls. All assays were performed with 6 replicates.

Chemo-surviving cell preparation

HCT-116 and HT-29 cells grown in monolayer were incubated with 50 μ M 5-FU and 1.25 μ M oxaliplatin (FOLFOX) (both 5-FU and oxaliplatin were obtained from Sigma-Aldrich, St. Louis, MO, USA) for 48 h, subsequently washed twice with phosphate-buffered saline (PBS). The cells that remained attached to the plate following the PBS wash (referred to as chemo-surviving cells) were trypsinized and regrown as a monolayer. Subsequent experiments of growth inhibition with gefitinib, curcumin and or FOLFOX were performed on these regrown chemo-surviving cells.

Western blot analysis

Western blot analysis was performed essentially according to our standard protocol (19). Briefly, the cells were solubilized in lysis buffer [50 mM Tris; 100 mM NaCl; 2.5 mM EDTA; 1% Triton X-100; 1% Nonidet P-40; 2.5 mM Na_3VO_4 ; 25 μ g/ml aprotinin; 25 μ g/ml leupeptin; 25 μ g/ml pepstatin A; and 1 mM phenylmethylsulfonyl fluoride (PMSF)]. Following clarification at 10,000 \times g for 15 min, the supernatant was used for Western blot analysis. In all analyses, protein concentration, determined by the Bio-Rad Protein Assay kit (Bio-Rad, Hercules, CA, USA), was standardized among the samples. Aliquots of cell lysates containing 50 μ g of protein were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Following electrophoresis, proteins were transferred electrophoretically onto supported nitrocellulose membranes (Osmonics, Gloucester, MA, USA). Membranes were incubated for 1 h at room temperature with blocking buffer, TBS-T (20 mM Tris, pH 7.6, 100 mM NaCl, 0.1% Tween-20) and 5% nonfat dry milk with gentle agitation. After washing the membranes with TBS-T, they were incubated overnight at 4°C in TBS-T buffer containing 5% milk and with one of the following antibodies (1:1000 dilution): phospho-EGFR (Tyr¹¹⁷³), phospho-ErbB-2/HER-2 (Tyr¹¹²¹), phospho-IGF-1R, phospho-Akt (Ser⁴⁷³) or COX-2 (Cell Signaling, Boston, MA, USA). The membranes were washed 3 times with TBS-T, and subsequently incubated with appropriate secondary antibodies (1:5000 dilutions) (Santacruz Biotechnology, Santa Cruz, CA, USA) in TBS-T containing 5% milk for 2 hours at room temperature with gentle agitation. The membranes were washed again with TBS-T, and the protein bands were visualized by enhanced chemiluminescence (ECL) detection system (Amersham, Sweden). The membranes containing the electrophoresed proteins were exposed to X-Omat film (PerkinElmer, Waltham, MA, USA). The membranes were then stripped (2 \times for 15 min at 55°C) in stripping buffer containing 100 mM 2-mercaptoethanol, 2% SDS and 62.5 mM Tris-HCl pH6.7, and re probed with β -actin as loading control. All Western blots were performed at least three times for each experiment. Densitometric measurements of the scanned bands were performed using the digitized scientific software program UN-SCNAT. Data were normalized to β -actin.

Transfection with si-RNAs

Chemo-surviving HCT-116 cells were transfected with the vector plasmid, EGFR si-RNA or IGF-1R si-RNA (Ambion, Austin, TX, USA) using Lipofectamine 2000 according to the manufacturer's instruction. For transfection, approximately 0.5–1 \times 10⁶ HT-29 or HCT-116 cells in 2 ml DMEM/10% FBS were plated in 96-well tissue culture dishes. After 24 h, the cells were transfected with Lipofectamine-2000 according to the manufacturer's instruction.

Two micrograms of vector, EGFR si-RNA or IGF-1R si-RNA were used in each well, and incubated for 48 h.

Statistical analysis

Data are represented as the mean±SEM. The statistical significance of differential findings between experimental groups and control was determined by Student's *t*-test as implemented by Excel 2000 (Microsoft Corp., Redmond, WA, USA). *P*-values smaller than 0.05 were considered statistically significant.

Results

An MTT assay was performed to determine the survival rate of colon cancer HCT-116 and HT-29 cells following FOLFOX treatment. We observed that 60–70% of the cells survived following 48 h incubation with FOLFOX (50 µM 5-FU and 1.25 µM oxaliplatin), henceforth referred to as chemo-surviving cells (Figure 1A data for HT-29 are not shown as they were similar to HCT-116).

Since members of EGFR family and IGF-1R play an important role in development and progression of colorectal cancer (5–8), we postulated that the chemo-surviving cells would exhibit increased expression and/or activation of survival promoting growth factor receptors, specifically EGFR and its family members and IGF-1R. Indeed, we observed a robust 2-fold and 4.8-fold increase in activation of EGFR and IGF-1R, respectively, in chemo-surviving HCT-116 cells (Figure 1B) compared to untreated cells (control). There was also a modest 30% increase in HER2 activation (Figure 1B). We also examined the expression and activation of AKT and COX-2, both of which are known to be critically involved in regulating cell survival signaling, downstream of growth factor receptors in various types of cancer including colorectal cancer. We observed a 20% increase in expression of COX-2 and a 78% increased activation of AKT without a significant change in its expression, when compared with the untreated parental cell controls (Figure 1B).

On the basis of our previous observation that curcumin synergizes with FOLFOX through down-regulation of EGFRs and IGF-1R, we hypothesized that curcumin would cause growth inhibition of chemo-surviving colon cancer cells. To test this hypothesis, chemo-surviving HCT-116 and HT-29 cells were subjected to further treatment for 48 h with FOLFOX or FOLFOX with two different doses of curcumin (25 or 50 µM). Incubation with the growth media only served as controls. Parental HCT-116 and HT-29 cells were also included to show changes in cell survival after 48 h FOLFOX treatment. Treatment of FOLFOX for 48 h resulted in approximately 50% and 60% survival of HCT-116 and HT-29 cells, respectively, when compared with the corresponding untreated parental cells. Further treatment of chemo-surviving HCT-116 and HT-29 cells with FOLFOX produced only a very modest growth inhibition (16–27%), when compared with the media-treated controls (Figure 2A). However, treatment of the chemo-surviving HCT-116 and HT-29 cells with FOLFOX together with either dose of curcumin exhibited significant growth inhibition compared to the media-treated controls (Figure 2A): low-dose curcumin (25 µM) in combination with FOLFOX was highly effective, producing a 60% growth inhibition in both HCT-116 and HT-29 chemo-surviving cells, and inhibition increased to 70–78% with the 50 µM dose of curcumin (Figure 2A).

Experiments were then performed to examine the effect of FOLFOX alone and in combination with curcumin (25 µM) on activation of the growth factor receptors and downstream mediators that are increased in chemo-surviving cells. Since curcumin was found to be effective in inhibiting growth of chemo-surviving colon cancer HCT-116 cells, whereas FOLFOX was not, we hypothesized that curcumin alone or in combination with

FOLFOX would result in attenuation of growth factor signaling that is upregulated in chemo-surviving cells. Indeed, continued treatment of chemo-surviving HCT-116 cells with FOLFOX resulted in either no change or only a slight increase in activation of EGFR, HER2 and IGF-1R as well as downstream signaling molecules AKT and COX-2 compared to the media-treated controls (Figure 2B). On the other hand, addition of curcumin to FOLFOX resulted in a marked 39–74% inhibition of activation of EGFR, HER2 and IGF-1R (Figure 2B). Additionally, there was a 91% reduction in the expression of COX-2 and a 36% attenuation in activation of AKT with the combined treatment of curcumin and FOLFOX, compared to the media-treated controls (Figure 2B).

Since EGFR activation is increased in chemo-surviving cells and is attenuated by curcumin, we sought to determine whether and to what extent the selective EGFR tyrosine kinase inhibitor gefitinib (Iressa) would be effective in growth inhibition of chemo-surviving HCT-116 colon cancer cells. Pretreatment of chemo-surviving HCT-116 cells with gefitinib (1 μ M) for 24 hours caused a 40% growth inhibition (Figure 3). The gefitinib-pretreated cells were then washed and subjected to treatment with the media alone (control), gefitinib, curcumin or curcumin plus FOLFOX for 48 h. Treatment with the media alone resulted in regrowth of chemo-surviving cells, whereas continued gefitinib treatment caused a 60% growth inhibition at the end of the 3 day experimental period (including a 24-h pretreatment period) (Figure 3). However, treatment with curcumin alone or curcumin and FOLFOX resulted in a marked >90% growth inhibition (Figure 3). It is noteworthy that addition of chemotherapy to curcumin resulted in no additional benefit over that observed with curcumin alone in inhibiting the growth of chemo-surviving colon cancer HCT-116 cells (Figure 3).

We then examined the effect of individual inhibition of EGFR or IGF-1R on growth inhibition of chemo-surviving HCT-116 cells. Transfection of chemosurviving HCT-116 cells with EGFR or IGF-1R si-RNA resulted in statistically significant 30% growth inhibition (Figure 4). Curcumin, which we have shown to inhibit EGFR as well as IGF-1R (19), caused a robust 70% growth inhibition at 48 h (Figure 4). Addition of curcumin to EGFR-si RNA and IGF-1R si-RNA transfected cells resulted in a minor further growth inhibition compared to curcumin alone (Figure 4).

Discussion

Chemotherapy produces incomplete responses in a vast majority of cancer cases particularly of colorectal cancer (2). This leads to survival of a population of cells within the tumor resulting in subsequent chemotherapy-resistant relapses. The precise mechanism of this phenomenon of chemo-survival remains unknown.

Abnormal activity of EGFRs has been associated with the development and progression of many malignancies, including that of the colon. In particular, overexpression of EGFR and HER2 in colorectal cancer correlates with an extremely poor clinical prognosis (20, 21). The majority of solid tumors, including those of the colon, overexpress one or more members of the EGFR family and coexpression of EGFR with HER-2 or HER-3 results in the development of enhanced drug resistance (9, 22). More recently, IGF-1R is also emerging as an important pathway responsible for the development and progression of colorectal cancer. In addition, there is crosstalk between EGFRs and IGF-1R, resulting in therapeutic resistance with targeting individual pathways (23). EGFR inhibitors have been successfully incorporated in the therapeutic armamentarium of colon cancer. However, the benefits appear to be modest and the complete responses are rare (24). One of the mechanisms of resistance to EGFR inhibitors is their heterodimerization with IGF-1R receptor (25). Hence, co-targeting of EGFR and IGF-1R would likely result in greater therapeutic efficacy. In our

model of chemo-surviving colon cancer cells, we show that not only EGFR and its family members, but also IGF-1R, are significantly activated, confirming their involvement in survival of these cells. Therefore, targeting them may provide beneficial effect in terms of reversing chemotherapy resistance. Recently a report by Dallas *et al.* who demonstrated inhibition of growth of oxaliplatin-resistant colon cancer cells by inhibiting IGF-1R, further supports the contention that certain growth factor receptors play a critical role in cells surviving chemotherapy insult (11). However, one of the major concerns with combining various biological agents is an increase in overall toxicity. Hence, the development of pleiotropic agents with minimal toxicity is highly desirable in combating the emergence of chemo-surviving cells and the resultant subsequent relapse. Our current data demonstrate that one can target these survival pathways with non-toxic pleiotropic agents such as curcumin for therapeutic gains.

Curcumin [diferuloylmethane; 1,7-*bis*-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione], the major active ingredient in turmeric powder, affects promotion/progression phases of colon carcinogenesis (12–16). The effect of curcumin on progression of the cancer is pleiotropic. A plethora of evidence suggests that the chemopreventive effect is mainly due to inhibition of COX-2, which leads to inhibition of prostaglandin E and of subsequent colon cancer growth (9, 26). COX-2 inhibition is due to attenuation of activation of transcription factors activator protein-1 (AP-1) and nuclear factor- κ B (NF- κ B), which regulate its transcription (26, 27). There is some evidence that curcumin inhibits the ligand-induced activation of EGFR and reduces its transcription *via* inhibition of *myc*-1 transcription factor (28). Moreover, our recent data demonstrate that curcumin inhibits both expression and activation of not only EGFR but also its family members and IGF-1R (19). In this report, we show that curcumin is not only highly effective in inhibiting the growth of chemo-surviving colon cancer cells but is also superior to EGFR inhibition by gefitinib and to IGF-1R attenuation through IGF-1R si-RNA. We believe that the ability of curcumin to effectively downregulate multiple growth factor receptor-mediated survival signals is critical in inhibiting chemo-surviving colon cancer cells. Curcumin has been shown to have no discernible toxicity in multiple human studies. Hence, it would be an ideal agent to use in combination with chemotherapy to prevent the emergence of chemo-surviving cells.

In conclusion, chemo-surviving colon cancer cells pose a major threat to the success of colon cancer chemotherapeutics *via* up-regulation of EGFRs and IGF-1R. Inclusion of curcumin, which effectively inhibits EGFRs and IGF-1R in chemo-surviving cells, into colon cancer chemotherapeutics is likely to inhibit the growth of the chemo-surviving cells and reduce the likelihood of subsequent relapse following chemotherapy at no additional toxicity.

Acknowledgments

The work was supported by grants from the National Institutes of Health/National Institute on Aging (APNM; R01 AG014343), Department of Veterans Affairs.

References

1. [last accessed on 11/30/06] <http://seer.cancer.gov/faststats/sites.php?site=Colon%20and%20Rectum%20Cancer&stat=Survival>
2. Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, Berlin J, Baron A, Griffing S, Holmgren E, Ferrara N, Fyfe G, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med.* 2004; 23:2335–2342. [PubMed: 15175435]
3. McCarty MF. Targeting multiple signaling pathway as a strategy for managing prostate cancer: multifocal signal modulation therapy. *Integ Cancer Ther.* 2004; 3:349–380.

4. Salomon DS, Brandt R, Ciardiello F, Normanno N. Epidermal growth factor-related peptides and their receptors in human malignancies. *Crit Rev Oncol Hematol*. 1995; 19:183–232. [PubMed: 7612182]
5. Messa C, Russo F, Caruso MG, Di Leo A. EGF, TGF- β and EGF-R in human colorectal adenocarcinoma. *Acta Oncol*. 1998; 37:285–289. [PubMed: 9677101]
6. Harari PM, Huang S-M. Modulation of molecular targets to enhance radiation. *Clin. Cancer Res*. 2000; 6:323–325. [PubMed: 10690506]
7. Hakam A, Yeatman TJ, Lu L, Mora L, Marcet G, Nicosia SV, Karl RC, Coppola D. Expression of insulin-like growth factor-1 receptor in human colorectal cancer. *Hum Pathol*. 1999; 30:1128–1133. [PubMed: 10534157]
8. Adachi Y, Lee CT, Coffee K, Yamagata N, Ohm JE, Park KH, Dikov MM, Nadaf SR, Arteaga CL, Carbone DP. Effects of genetic blockade of insulin-like growth factor receptor in human colon cancer cell lines. *Gastroenterology*. 2002; 123:1191–1204. [PubMed: 12360481]
9. Chen X, Yeung TK, Wang Z. Enhanced drug resistance in cells coexpressing ErbB2 with EGF receptor or ErbB3. *Biochem Biophys Res Commun*. 2000; 277:757–763. [PubMed: 11062025]
10. Dixit M, Yang JL, Poirier MC, Price JO, Andrews PA, Arteaga CL. Abrogation of cisplatin-induced programmed cell death in human breast cancer cells by epidermal growth factor antisense RNA. *J Natl Cancer Inst*. 1997; 89:365–373. [PubMed: 9060958]
11. Dallas NA, Xia L, Fan F, Gray MJ, Gaur P, van Buren G 2nd, Samuel S, Kim MP, Lim SJ, Ellis LM. Chemoresistant colorectal cancer cells, the cancer stem cell phenotype, and increased sensitivity to insulin-like growth factor-I receptor inhibition. *Cancer Res*. 2009; 69:1951–1957. [PubMed: 19244128]
12. Hanif R, Qiao L, Shiff SJ, Rigas B. Curcumin, a natural plant phenolic food additive, inhibits cell proliferation and induces cell cycle changes in colon adenocarcinoma cell lines by a prostaglandin-independent pathway. *J Lab Clin Med*. 1997; 130:576–584. [PubMed: 9422331]
13. Chauhan DP. Chemotherapeutic potential of curcumin for colorectal cancer. *Curr Pharm Des*. 2002; 8:1695–1706. [PubMed: 12171541]
14. Huang MT, Wang ZY, Georgiadis CA, Laskin JD, Conney AH. Inhibitory effects of curcumin on tumor initiation by benzo[a]pyrene and 7,12-dimethylbenz[a]anthracene. *Carcinogenesis*. 1992; 13:2183–2186. [PubMed: 1423891]
15. Rao CV, Simi B, Reddy BS. Inhibition by dietary curcumin of azoxymethane-induced ornithine decarboxylase, tyrosine protein kinase, arachidonic acid metabolism and aberrant crypt foci formation in the rat colon. *Carcinogenesis*. 1993; 14:2219–2225. [PubMed: 8242846]
16. Rao CV, Rivenson, Simi B, Reddy BS. Chemoprevention of colon carcinogenesis by dietary curcumin, a naturally occurring plant phenolic compound. *Cancer Res*. 1995; 55:259–266. [PubMed: 7812955]
17. Perkins S, Verschoyle RD, Hill K, Parveen I, Threadgill MD, Sharma RA, Williams ML, Steward WP, Gescher AJ. Chemopreventive efficacy and pharmacokinetics of curcumin in the Min/+ mouse, a model of familial adenomatous polyposis. *Cancer Epidem Biomark Prev*. 2002; 11:535–540.
18. Sharma RA, Euden SA, Platton SL, Cooke DN, Shafayat A, Hewitt HR, Marczyklo TH, Morgan B, Hemingway D, Plummer SM, Pirmohamed M, Gescher AJ, et al. Phase I clinical trial of oral curcumin: biomarkers of systemic activity and compliance. *Clin Cancer Res*. 2004; 10:6847–6854. [PubMed: 15501961]
19. Patel BB, Sengupta R, Qazi S, Vachhani H, Yu Y, Rishi AK, Majumdar AP. Curcumin enhances the effects of 5-fluorouracil and oxaliplatin in mediating growth inhibition of colon cancer cells by modulating EGFR and IGF-1R. *Int J Cancer*. 2008; 122:267–273. [PubMed: 17918158]
20. Galizia G, Lieto E, Ferraraccio F, De Vita F, Castellano P, Orditura M, Imperatore V, La Mura A, La Manna G, Pinto M, Catalano G, Pignatelli C, et al. Prognostic significance of epidermal growth factor receptor expression in colon cancer patients undergoing curative surgery. *Ann Surg Oncol*. 2006; 6:823–835. [PubMed: 16614884]
21. Park DI, Kang MS, Oh SJ, Kim HJ, Cho YK, Sohn CI, Jeon WK, Kim BI, Han WK, Kim H, Ryu SH, Sepulveda AR. HER-2/neu overexpression is an independent prognostic factor in colorectal cancer. *Int J Colorectal Dis*. 2006; 5:491–497. [PubMed: 16947041]

22. Pinkas-Kramarski R, Soussan L, Waterman H, Levkowitz G, Alroy I, Klapper L, Lavi S, Seger R, Ratzkin BJ, Sela M, Yarden Y. Diversification of Neu differentiation factor and epidermal growth factor signaling by combinatorial receptor interactions. *EMBO J.* 1996; 15:2452–2467. [PubMed: 8665853]
23. Jin Q, Esteva FJ. Cross-talk between the ErbB/HER family and the type I insulin-like growth factor receptor signaling pathway in breast cancer. *J Mammary Gland Biol Neoplasia.* 2008; 13:485–98. Review. [PubMed: 19034632]
24. Jonker DJ, O’Callaghan CJ, Karapetis CS, et al. Cetuximab for the treatment of colorectal cancer. *N Engl J Med.* 2007; 357:2040–2048. [PubMed: 18003960]
25. Morgillo F, Woo JK, Kim ES, Hong WK, Lee HY. Heterodimerization of insulin-like growth factor receptor/ epidermal growth factor receptor and induction of survivin expression counteract the antitumor action of erlotinib. *Cancer Res.* 2006; 66:10100–10111. [PubMed: 17047074]
26. Kawamori T, Lubet R, Steele VE, Kelloff GJ, Kaskey RB, Rao CV, Reddy BS. Chemopreventive effect of curcumin, a naturally occurring anti-inflammatory agent, during the promotion/ progression stages of colon cancer. *Cancer Res.* 1991; 59:597–601. [PubMed: 9973206]
27. Plummer SM, Holloway KA, Manson MM, Munks RJ, Kaptein A, Farrow S, Howells L. Inhibition of cyclo-oxygenase-2 expression in colon cells by the chemopreventive agent curcumin involves inhibition of NF- κ B activation *via* the NIK/IKK signaling complex. *Oncogene.* 1999; 18:6013–6020. [PubMed: 10557090]
28. Chen A, Xu J, Johnson AC. Curcumin inhibits human colon cancer cell growth by suppressing gene expression of epidermal growth factor receptor through reducing the activity of the transcription factor Egr-1. *Oncogene.* 2006; 2:278–287. [PubMed: 16170359]




Figure 1.
 A: Proportion of chemo-surviving HCT-116 cells following 48 h of treatment with FOLFOX (50 μ M 5-FU and 1.25 μ M oxaliplatin) chemotherapy, as measured by MTT assay. B: Western-blot showing the levels of activated (tyrosine phosphorylated) forms of EGFR, HER-2, IGF-1R, and downstream mediators AKT, as well as COX-2, in chemo-surviving HCT-116 cells.




Figure 2.

A: Growth inhibition (MTT assay) of chemo-surviving HCT-116 and HT-29 cells (pre treated with FOLFOX for 48 hours) with FOLFOX alone or in combination with curcumin (25 and 50 µM), compared to media-treated cells (control) following additional 48 h treatment. Untreated parental cells are included for purposes of comparison (total of 96 hours' exposure to media alone). B: Western blot showing the levels of activated growth factor receptors EGFR, HER-2, IGF-1R, and downstream mediators AKT, as well as COX-2, in chemo-surviving HCT-116 cells following treatment with FOLFOX alone or in combination with curcumin (25 and 50 µM), compared to media-treated cells (control). Parental untreated cells were included for purposes of comparison (total of 96 hours exposure to media alone). *p<0.001 Compared to control.




Figure 3. Growth inhibition (MTT assay) of chemo-surviving HCT-116 cells pretreated with 1 μM gefitinib followed by additional 48 h treatment with gefitinib, curcumin or FOLFOX and curcumin compared to cells in media alone (control).




Figure 4. Growth inhibition (MTT assay) of chemo-surviving HCT-116 cells transfected with vector, EGFR si-RNA, IGF-1R si-RNA and treated with either DMSO (control) or curcumin. * $p < 0.01$ and ** $p < 0.001$ compared to control.